


DERIVATIVES OF THE MESODERMAL GERM LAYER

Paraxial mesoderm




It develops into
TWO PERIPHERAL MASSES
and a **constriction in the middle**

Called:

1-Medial mesoderm

2-Intermediate mesoderm

3-lateral mesoderm




Medial mesoderm

- The medial mesoderm enlarges pushing the ectoderm upwards to give the **somites**

➤ As the embryo develops the number of the somites increases from one to reach about 44-45 somites

➤ when the embryo is completely developed

About 10 somites vanish when the tail of the embryo is lost




- ❖ **The first pair** of somites arises in the **occipital region** of the embryo at approximately the 20th day of development
- ❖ From here, new somites appear in craniocaudal sequence at a rate of approximately three pairs per day until the end **of the fifth week**,



There are:
four occipital
eight cervical
12 thoracic
five lumbar
five sacral,
and eight to 10 coccygeal pairs.

The
first occipital and the last five to seven coccygeal somites later disappear


WHAT IS THE destiny OF EACH SOMITE?



By the beginning of the fourth week
cells forming the *ventral and medial walls of the somite*
lose their compact organization,
and shift their position to surround the notochord
These cells, collectively known as

THE SCLEROTOME

They will
surround the
spinal cord and
notochord to form
**the
vertebral
column**



Cells at the dorsolateral portion of the somite also migrate as precursors of the

limb and body wall

musculature



(hypomeric) musculature

After migration of these muscle cells and cells of the sclerotome,

Cells at the dorsomedial portion of the somite proliferate and migrate to form a new layer



THE MYOTOME

myotome contributes to muscles of the back (epaxial musculature)
or epimeric musculature
the **extensor muscles of the vertebral column**






*The remaining dorsal
epithelium
forms the dermatome*

dermatomes form
the dermis and subcutaneous tissue of the skin



a transversal section through the embryo at level A is displayed. The somites have released themselves and form *dermatomes*, *myotomes* and *sclerotomes*.






DERIVATIVES OF THE INTERMEDIATE MESODERM

It gives off:

**1- Urine performing tubule (Kidney
and ureter)**

**2-internal genitalia in males and femals
(part of it not all)**



Embryonic folds

WHY the embryo folds?


1- Extensive and rapid growth of the cranial end of the neural tube

2- The faster growth of the axial part of the embryonic disc than its periphery

3- Enlargement of the amnion

Folding of the embryo Cephalocaudally and Laterally

Head (cephalic) fold




Tail (caudal) fold


The embryonic disc begins to bulge into the amniotic cavity and to fold

Cephalocaudally


Folding of the embryo Cephalocaudally




Folding of the embryo Laterally



DERIVATIVES OF
THE LATERAL
MESODERM




The intraembryonic coelom (body cavity) first appears as small, isolated, **coelomic spaces in the lateral intraembryonic mesoderm**



These spaces coalesce to form a single, horseshoe-shaped cavity—the **intraembryonic coelom**

The coelom divides the lateral mesoderm into two layers:




Lateral mesoderm splits into two layers:

1- Parietal (somatic)

2- Visceral (splanchnic)

➤ Mesoderm from **the parietal layer**, together with **overlying ectoderm**, forms **the lateral body wall folds**

➤ These folds, together with the head (cephalic) and tail (caudal) folds, **close the ventral body wall**



1-The parietal layer of lateral mesoderm forms:


A) The dermis of the skin in the body wall and limbs

B) The bones and connective tissue of the limbs

C) The sternum

D) Mesoderm cells of the parietal layer surrounding the intraembryonic cavity form thin membranes, the mesothelial membranes, or serous membranes, which will line the
1-peritoneal
2- pleural 3- pericardial cavities


E) In addition, sclerotome and muscle precursor cells that migrate into the parietal layer of lateral plate mesoderm form
the costal cartilages,
limb muscles,
and most of the body wall muscles



2-The visceral layer of lateral mesoderm

Surrounds the primitive gut and together with embryonic endoderm, forms

THE WALL
OF THE
GUT TUBE



Mesoderm also gives rise to the **vascular system, that is, the heart, arteries, veins, lymph vessels, and all blood and lymph cells**





Figure 5.15 Extraembryonic blood vessel formation in the villi, chorion, connecting stalk, and wall of the yolk sac in a presomite embryo of approximately 19 days.

**DERIVATIVES OF
THE ENDODERMAL GERM
LAYER**


➤ The gastrointestinal tract is the main organ system derived from the endodermal germ layer

➤ *With development the embryonic disc begins to bulge into the amniotic cavity and to fold cephalocaudally and Lateral folds also form and move ventrally to assist in body wall closure*





A

As a result of cephalocaudal folding, a continuously larger portion of the endodermal germ layer is incorporated into the body of the embryo to form the gut tube.



The midgut communicates with the yolk sac **by way of a broad stalk the VITELLINE DUCT**



The tube is divided into three regions:

FOREGUT

MIDGUT

HINDGUT


A


The midgut remains in communication with the yolk sac.

Initially, this connection is wide but as a result of body folding, it gradually becomes long and narrow to form

the vitelline duct

Only much later, when the vitelline duct is obliterated, does the midgut lose its connection with the original endoderm-lined cavity and obtain its free position in the abdominal cavity





At its cephalic end, **the foregut** is temporarily bounded by


an ectodermal-endodermal
(no mesoderm)

membrane called the

OROPHARYNGEAL membrane

This membrane separates the stomodeum, (the primitive oral cavity derived from ectoderm), from the pharynx, (a part of the foregut derived from endoderm).

In the fourth week, the oropharyngeal membrane ruptures, establishing an open connection between **the oral cavity** and the primitive gut



The hindgut also terminates temporarily at an ectodermal-endodermal membrane,

THE CLOACAL membrane

This membrane separates the upper part of the anal canal (derived from endoderm), from the lower part called (the proctoderm) that is formed by an invaginating pit lined by ectoderm.


The membrane breaks down in the seventh week to create the opening for the **anus**

No mesoderm

**Buccopharyngeal
membrane**

**Cloacal
membrane**

WHY?



**As a result of folding from the head, tail, and
two lateral body wall folds**

**The ventral body wall of the embryo is closed
except for a small part in the umbilical region
where the yolk sac duct and connecting stalk
are attached.**

ENDODERM GIVES RISE TO:

- The epithelial lining of the respiratory tract
- The parenchyma of the thyroid, parathyroids, liver, and pancreas
- The reticular stroma of the tonsils and thymus
- The epithelial lining of the urinary bladder and urethra
- The epithelial lining of the tympanic cavity and auditory tube