

## Carbohydrates Metabolism



Review of Carbohydrates  
Digestion<sup>1</sup> and absorption<sup>2</sup> of carbohydrates

Suggested Readings:


- 1: Lippincot's Illustrated reviews: Biochemistry
- 2: Marks' Basic Medical Biochemistry

## Carbohydrates Metabolism

- Objectives
  - Utilization of Glucose → Energy
  - Non-Carbohydrates → Glucose
  - Storage of Glucose → Glycogen
  - Release of Glucose from Glycogen
  - Reducing Power NADPH >> GSH
  - Glucuronic acid >> Drug metabolism
  - Interconversion of sugars







## Examples of an aldose and ketose




## Examples of monosaccharides found in human

<u>Generic names</u>	<u>Examples</u>
<b>3</b> carbons: trioses	Glyceraldehyde
<b>4</b> carbons: tetroses	Erythrose
<b>5</b> carbons: pentoses	Ribose
<b>6</b> carbons: hexoses	Glucose
<b>7</b> carbons: heptoses	Sedoheptulose
<b>9</b> carbons: nonoses	Neuraminic acid












Disaccharide: A sugar made of two sugar units joined by glycosidic bond

Maltose: a disaccharide made from two glucose units










## Mucosal cell membrane-bound enzymes


ENZYME	Bond Cleaved	Substrates
Isomaltase	$\alpha$ 1 $\rightarrow$ 6	Isomaltose
Maltase	$\alpha$ 1 $\rightarrow$ 4	Maltose
Sucrase	$\alpha$ 1 $\rightarrow$ 2	Sucrose
Lactase	$\beta$ 1 $\rightarrow$ 4	Lactose
Trehalase	$\alpha$ 1 $\rightarrow$ 1	Trehalose
Exoglucosidase	$\alpha$ 1 $\rightarrow$ 4	Glucoamylose



**FIG. 27.5.** The major portion of the sucrase–isomaltase complex, containing the catalytic sites, protrudes from the absorptive cells into the lumen of the intestine. Other domains of the protein form a connecting segment (stalk) and an anchoring segment that extends through the membrane into the cell. The complex is synthesized as a single polypeptide chain that is split into its two enzyme subunits extracellularly. Each subunit is a domain with a catalytic site (distinct sucrase–maltase and isomaltase–maltase sites). In spite of their maltase activity, these catalytic sites are often called just *sucrase* and *isomaltase*.

## Abnormal Degradation of disaccharides

- Lactase deficiency:
  - ½ world's population
- Sucrase isomaltase deficiency:
- Causes:
  - Genetics
  - Variety of intestinal diseases
  - Malnutrition
  - Injury of mucosa ie by drugs
  - Severe diarrhea



## Absorption of Sugars


Polar molecules can not diffuse

**A:** Na<sup>+</sup>-independent facilitated diffusion transport

GLUT 1-----GLUT 14

Glc. Movement follows concentration gradient


Two conformation states



## Na<sup>+</sup> monosaccharide cotransporter system (SGLT)

- \* Against concentration gradient.
- \* Small intestine: Active uptake from lumen of intestine.
- \* Kidney: reabsorption of glucose in proximal tubule.

## Na<sup>+</sup> monosaccharide cotransporter system (SGLT)



**Table 27.5 Properties of the GLUT 1 to GLUT 5 Isoforms of the Glucose Transporter Proteins**

Transporter	Tissue Distribution	Comments
GLUT 1	Human erythrocyte Blood-brain barrier Blood-retinal barrier Blood-placental barrier Blood-testis barrier	Expressed in cell types with barrier functions; a high-affinity glucose transport system
GLUT 2	Liver Kidney Pancreatic $\beta$ -cell Serosal surface of intestinal mucosa cells	A high-capacity, low-affinity transporter May be used as the glucose sensor in the pancreas
GLUT 3	Brain (neurons)	Major transporter in the central nervous system; a high-affinity system
GLUT 4	Adipose tissue Skeletal muscle Heart muscle	Insulin-sensitive transporter. In the presence of insulin, the number of GLUT 4 transporters increases on the cell surface; a high-affinity system
GLUT 5	Intestinal epithelium Spermatozoa	This is actually a fructose transporter
GLUT 7	Glucogenic tissues	at endoplasmic reticulum membrane

